arXiv:1304.5866v1 [math.CA] 22 Apr 2013

SOME REMARKS ON DUNKL-TYPE OPERATORS WITH PROJECTIONS TERMS

FETHI BOUZEFFOUR

Abstract. We present a Dunkl-type operator T_ξ, ($\xi \in \mathbb{R}^N$) constructed with the help of orthogonal projections associated with a subsystem R of root system. Similarly to the Dunkl theory we prove in the case when R is orthogonal system, the operators T_ξ, $\xi \in \mathbb{R}^N$ commute, also we construct an intertwining between T_ξ and the directional derivative ∂_ξ. In the one variable we prove that this operator admits as eigenfunction the Kummer function. This paper contains some other results which are the prelude of harmonic analysis related to this operator.

Keywords: Root system, Differential-difference operator, intertwining operator.

1. Introduction

During the last thirty years, Dunkl operators have largely contributed to the development of the harmonic analysis and the theory of special functions associated with root systems. Roughly speaking, Dunkl operators are commuting differential-difference operators on a Euclidean space which are associated with a finite refection group and have a continuous set of parameters, called the multiplicities.

The Dunkl operators were introduced by C.F. Dunkl in a series of papers (see, [3, 4, 5, 6]), where he built up the framework for a theory of special functions and integral transforms in several variables related with reflection groups. Besides them, there are now various further Dunkl-type operators, in particular the trigonometric Dunkl operators of Heckman (see,[9, 10]), Opdam (see, [11]) and Cherednik (see, [2]) and the important q-analogues of Macdonald and Cherednik (see, [8]).

In this work, we introduce a new class of Dunkl-type operators T_ξ, $\xi \in \mathbb{R}^N$, which are a mixture of differential and difference operators with orthogonal projections terms. In other words, these operators flow from Dunkl operators after replacing the reflections terms that exist in their expressions by orthogonal projections terms.

Several problems related to the Dunkl theory arise in the sitting of ours operators, in particular, commutativity of T_ξ, $\xi \in \mathbb{R}^N$, the existence of intertwining operator.

In this paper, we show that these operators satisfy the fundamental property of differential-difference operators that is their commutativity in particular case when the subsystem R is orthogonal and we construct an intertwining between T_ξ and the directional derivative. The specialization of this theory to the case of rank one has its own interest, because everything can be done there in a much more explicit way, and new results for special functions in one variable can be obtained. In the rank one case, the eigenfunction function associated to the Dunkl-type operator is expressed explicitly in terms of the Kummer function.

The outline of the content of this paper is as follows. In section 2, we collect some definitions and results related root systems and Dunkl operators which will be relevant for the sequel, also we introduce a Dunkl-type operators T_ξ and we show that these operators commute in the case of orthogonal system. In section 3, we give an explicit formula for the intertwining operator between T_ξ and the directional derivative. Section 4, is devoted to study two interesting case, the one variable case and the case when the R is the canonical basis of $\mathbb{R}^N$.

2. The operator T_ξ

Let us begin to recall some results concerning the root systems. A useful reference for this topic is the book by Humphreys [7]. Let $\alpha \in \mathbb{R}^N\backslash\{0\}$, we denote by s_α the reflection in the hyperplane $(\mathbb{R}\alpha)^\perp$ orthogonal to α; that is,

$$s_\alpha(x) = x - 2\frac{\langle x, \alpha\rangle}{|\alpha|^2}\alpha,$$

where $\langle .\, , .\rangle$ denotes the Euclidean scalar product on $\mathbb{R}^N$, and $|x| = \sqrt{\langle x, x\rangle}$.
A root system is a finite set R of nonzero vectors in $\mathbb{R}^N$ such that

$$\sigma_\alpha(R) = R, \qquad \text{and} \qquad R \cap \mathbb{R}\alpha = \{\pm\alpha\}$$

for any $\alpha \in R$. A positive subsystem R_+ is any subset of R satisfying $R = R_+ \cup \{-R_+\}$. The real Coxeter group $W = W(R)$ (or real finite reflection group) generated by the root system $R \subset \mathbb{R}^N$ is the subgroup of orthogonal group $O(N)$ generated by $\{s_\alpha : \alpha \in R\}$.
A multiplicity function on R is a complex-valued function $\kappa : R \longrightarrow \mathbb{C}$ which is invariant under the Coxeter group, i.e.,

$$\kappa(\alpha) = \kappa(g\alpha), \qquad \forall\ \alpha \in R, \quad \forall\ g \in W.$$

Let $\xi \in \mathbb{R}^N$, the Dunkl operator T_ξ, associated with the Coxeter group $W(R)$ and the multiplicity function κ, is the first order differential-difference operator:

$$(T_\xi f)(x) \ = \ \partial_\xi f(x) + \sum_{\alpha \in R_+} \kappa(\alpha)\langle \alpha, \xi\rangle \frac{f(x) - f(s_\alpha x)}{\langle x, \alpha\rangle}. \tag{2.1}$$

Here ∂_ξ is the direction derivative corresponding to ξ.
The Dunkl operator T_ξ is a homogeneous differential operator of degree -1. By the W-invariance of the multiplicity function κ, we have

$$g^{-1} \circ T_\xi \circ g = T_{g\xi}, \qquad \forall\ g \in W(R), \quad \xi \in \mathbb{R}^N.$$

The remarkable property of the Dunkl operators is that the family $\{T_\xi, \xi \in \mathbb{R}^N\}$ generates a commutative algebra of linear operators on the $\mathbb{C}$-algebra of polynomial functions.
For a vector $\alpha \in \mathbb{R}^N \setminus \{0\}$, we write

$$\tau_\alpha(x) = x - \frac{\langle x, \alpha\rangle}{|\alpha|^2}\alpha, \ x \in \mathbb{R}^N, \tag{2.2}$$

for the orthogonal projection in the hyperplane $(\mathbb{R}\alpha)^\perp = \{x : \langle x, \alpha\rangle = 0\}$, so that the reflection s_α with respect to hyperplane orthogonal to α is related to τ_α by

$$\tau_\alpha = \frac{1}{2}(1 + s_\alpha).$$

The hyperplane $(\mathbb{R}\alpha)^\perp$ is the invariant set of τ_α. Tow orthogonal projections τ_α and τ_β commute if and only if $\langle \alpha, \beta\rangle = 0$. The conjugate of orthogonal projection in a hyperplane is again an orthogonal projection in a hyperplane: suppose $u \in O(N)$ and $\alpha \in \mathbb{R}^N\backslash\{0\}$ then

$$u\tau_\alpha u^{-1} = \tau_{u\alpha}.$$

For $\xi \in \mathbb{R}^N\backslash\{0\}$, introduce the operator ρ_α defined on $C^\infty(\mathbb{R}^N)$ by

$$(\rho_\alpha f)(x) = \frac{f(x) - f(\tau_\alpha x)}{\langle x, \alpha\rangle}. \tag{2.3}$$

We denote by Π^N the $\mathbb{C}-$algebra of polynomials functions on $\mathbb{R}^N$ and Π^N_n the subspace of homogenous polynomials of degree n.

Lemma 2.1. *1) If $f \in C^\infty(\mathbb{R}^N)$, then $\rho_\alpha f \in C^\infty(\mathbb{R}^N)$.*
2)The operator ρ_α is a homogeneous difference operator of degree -1 on Π^N, that is, $\rho_\alpha f \in \Pi^N_{n-1}$, for $f \in \Pi^N_n$.

Proof. If $x \in \mathbb{R}^N$, then

$$(\rho_\alpha f)(x) = \frac{f(x) - f(\tau_\alpha x)}{\langle x, \alpha\rangle} = \frac{1}{|\alpha|^2}\int_0^1 \partial_\alpha f(x - t\frac{\langle x, \alpha\rangle}{|\alpha|^2}\alpha)\, dt. \tag{2.4}$$

The right hand side is smooth if f is, which proves the statement about $C^\infty(\mathbb{R}^N)$. If f is a polynomial, so is $\rho_\alpha f$. □

Lemma 2.2. *For $\alpha \in \mathbb{R}^N\backslash\{0\}$ and $f \in C^\infty(\mathbb{R})$. we have*

$$(\rho_\alpha f)(\tau_\alpha x) = |\alpha|^{-2}(\partial_\alpha f)(\tau_\alpha x).$$

Proof. Let $f \in C^\infty(\mathbb{R})$. From (2.4), we have

$$\begin{aligned}(\rho_\alpha f)(\tau_\alpha x) &= \frac{1}{|\alpha|^2}\int_0^1 \partial_\alpha f(\tau_\alpha x - t\frac{\langle \tau_\alpha x, \alpha\rangle}{|\alpha|^2}\alpha)dt\\ &= \frac{1}{|\alpha|^2}\int_0^1 \partial_\alpha f(\tau_\alpha x - t\frac{\langle x, \tau_\alpha \alpha\rangle}{|\alpha|^2}\alpha)dt\\ &= |\alpha|^{-2}(\partial_\alpha f)(\tau_\alpha x).\end{aligned}$$

□

Let R be a root system in $\mathbb{R}^N$, G the associated refection group and κ a multiplicity function on R. For $\xi \in \mathbb{R}^N$, we define T_ξ the differential-difference operator with orthogonal projections terms associated with R and κ by

$$(T_\xi f)(x) \;=\; \partial_\xi f(x) + \sum_{\alpha\in R}\kappa(\alpha)\langle\alpha, \xi\rangle\frac{f(x) - f(\tau_\alpha x)}{\langle x, \alpha\rangle}. \tag{2.5}$$

For $j = 1$, ..., N denote T_{e_j} by T_j. The above definition does not depend on the special choice of R. The Dunkl-type operators T_ξ attached with R can be considered as deformations of the usual partial derivatives by projections parts. These projections terms are coupled by parameters, which are given in terms of a multiplicity function κ. In case $\kappa = 0$, the operator T_ξ reduces to the corresponding directional derivative.

Example 2.1. In the rank one case, the root system is of type A_1 and the corresponding reflection s and orthogonal projection τ are given by

$$s(x) = -x, \;\; \tau(x) = \frac{1}{2}(1 + s)(x) = 0.$$

The Dunkl-type operator T_κ associated with the projection τ and the multiplicity parameter κ, ($\kappa \in \mathbb{C}$) is given by

$$\begin{aligned}T_\kappa f(x) &= f'(x) + \kappa\frac{f(x) - f(\tau(x))}{x}\\ &= f'(x) + \kappa\frac{f(x) - f(0)}{x}.\end{aligned}$$

Proposition 2.3. *1) If $f \in C^m(\mathbb{R}^N)$ with $m \geq 1$, then $T_\xi f \in C^{m-1}(\mathbb{R}^N)$.*
2)The operator T_ξ is a homogeneous differential-difference operator of degree -1 on Π^N, that is, $T_\xi p \in \Pi^N_{n-1}$, for $p \in \Pi^N_n$.

Consider the natural action of G on function $f : \mathbb{R}^N \to \mathbb{C}$ given by

$$(\sigma f)(\sigma^{-1}x),\ \sigma \in G.$$

Proposition 2.4. *Let $w \in G$ and $\xi \in \mathbb{R}^N$, then $wT_\xi w^{-1} = T_{w\xi}$.*

Proof. Let $f \in C^\infty(\mathbb{R})$, then

$$\begin{aligned} wT_\xi w^{-1} f\,(x) &= \partial_{w\xi} f(x) + w\sum_{\alpha\in R}\kappa(\alpha)\langle\alpha,\xi\rangle\frac{f(wx)-f(\tau_{w\alpha}x)}{\langle x,\alpha\rangle} \\ &= \partial_{w\xi} f(x) + \sum_{\alpha\in R}\kappa(\alpha)\langle\alpha,\xi\rangle\frac{f(x)-f(w\tau_\alpha w^{-1}x)}{\langle w^{-1}x,\alpha\rangle} \\ &= \partial_{w\xi} f(x) + \sum_{\alpha\in R}\kappa(\alpha)\langle\alpha,\xi\rangle\frac{f(x)-f(\tau_{w\alpha}x)}{\langle x,w\alpha\rangle} \\ &= T_{w\xi} f\,(x)\,. \end{aligned}$$

The summation variable $\alpha \in R$ is replaced by $\beta = w\alpha$. ☐

For $t \in \mathbb{R}^N$ let m_t denote the multiplier operator on Π^N given by

$$(m_t p)\,(x) := \langle t,x\rangle\, p\,(x)\,.$$

Proposition 2.5. *For $\alpha, t \in \mathbb{R}^N$*

$$[T_\xi, m_t]\, f(x) = \langle\xi,t\rangle\, f(x) + \sum_{\alpha\in R}\kappa(\alpha)\frac{\langle\xi,\alpha\rangle\langle t,\alpha\rangle}{|\alpha|^2} f(\tau_\alpha x),$$

where the commutator of two operators A, B is defined by

$$[A,B] := AB - BA$$

.

Proof. Let $f \in \Pi^N$, we have

$$\partial_\xi(\langle t,\,.\rangle\, f)\,(x) - \langle t,x\rangle\,\partial_\xi f(x) = \langle t,\xi\rangle\, f\,(x)\,.$$

Next for any $\alpha \in R$ we have

$$\frac{\langle t,x\rangle\, f(x) - \langle t,\tau_\alpha x\rangle\, f(\tau_\alpha x)}{\langle x,\alpha\rangle} - \langle t,x\rangle\,\frac{f(x)-f(\tau_\alpha x)}{\langle x,\alpha\rangle} = \frac{\langle t,\alpha\rangle}{|\alpha|^2} f(\tau_\alpha x).$$

This proves the formula. ☐

If τ_α is an orthogonal projection, then τ_α acts on function $f : \mathbb{R}^N \to \mathbb{C}$ as

$$(\tau_\alpha f)(x) := f(\tau_\alpha x).$$

Lemma 2.6. *For $\xi \in \mathbb{R}^N$ and $\beta \in R$ satisfying*

$$\forall\alpha \in R\backslash\{\beta\}\ \tau_\beta(\alpha) = \alpha.$$

Then

$$[\tau_\beta, T_\xi] = (1+\kappa(\beta))\frac{\langle\xi,\beta\rangle}{|\beta|^2}\tau_\beta\partial_\beta.$$

Proof. A simple calculation shows

$$(T_\xi \tau_\beta f)(x) = \partial_{\tau_\beta \xi} f(\tau_\beta x) + \sum_{\alpha\in R} \kappa(\alpha)\langle \alpha, \xi\rangle \frac{f(\tau_\beta x) - f(\tau_\beta \tau_\alpha x)}{\langle x, \alpha\rangle}.$$

Using Lemma 2.2 and the fact that $\tau_\beta(\alpha) = \alpha$, we can write

$$\begin{aligned}(\tau_\beta T_\xi f)(x) &= \partial_\xi f(\tau_\beta x) + \sum_{\alpha\in R\backslash\{\beta\}} \kappa(\alpha)\langle \alpha, \xi\rangle \frac{f(\tau_\beta x) - f(\tau_\alpha \tau_\beta x)}{\langle \tau_\beta x, \alpha\rangle} + \kappa(\beta)\frac{\langle \xi, \beta\rangle}{|\beta|^2}\tau_\beta \partial_\beta \\ &= \partial_\xi f(\tau_\beta x) + \sum_{\alpha\in R\backslash\{\beta\}} \kappa(\alpha)\langle \alpha, \xi\rangle \frac{f(\tau_\beta x) - f(\tau_\alpha \tau_\beta x)}{\langle x, \alpha\rangle} + \kappa(\beta)\frac{\langle \xi, \beta\rangle}{|\beta|^2}\tau_\beta \partial_\beta.\end{aligned}$$

On the other hand

$$(T_\xi \tau_\beta f)(x) = +\sum_{\alpha\in R} \kappa(\alpha)\langle \alpha, \xi\rangle \frac{f(\tau_\beta x) - f(\tau_\beta \tau_\alpha x)}{\langle x, \alpha\rangle},$$

Therefore,

$$\begin{aligned}[\tau_\beta, T_\xi] &= \partial_{\xi - \tau_\beta \xi} f(\tau_\beta x) + \kappa(\beta)\frac{\langle \xi, \beta\rangle}{|\beta|^2}\tau_\beta \partial_\beta \\ &= (1 + \kappa(\beta))\frac{\langle \xi, \beta\rangle}{|\beta|^2}\tau_\beta \partial_\beta.\end{aligned}$$

□

The main result about Dunk-type operators T_ξ, on which all their applications are based, is the following theorem

Theorem 2.7. *Let R be an orthogonal subsystem of a root system and $\xi, \eta \in \mathbb{R}^N$ then*

$$[T_\xi, T_\eta] = 0.$$

Proof. Let $t \in \mathbb{R}^N$; we will show $[[T_\xi, T_\eta], m_t] = 0$. Indeed

$$[[T_\xi, T_\eta], m_t] = [[T_\xi, m_t], T_\eta] - [[T_\eta, m_t], T_\xi],$$

and

$$\begin{aligned}[[T_\xi, m_t], T_\eta] &= [\langle \xi, t\rangle, T_\eta] + \sum_{\alpha\in R_+} \kappa(\alpha)\frac{\langle \xi, \alpha\rangle\langle t, \alpha\rangle}{|\alpha|^2}[\tau_\alpha, T_\eta] \\ &= \sum_{\alpha\in R_+} (1 + \kappa(\alpha))\kappa(\alpha)\frac{\langle \xi, \alpha\rangle\langle t, \alpha\rangle\langle \eta, \alpha\rangle}{|\alpha|^4}\tau_\alpha \partial_\alpha,\end{aligned}$$

which is symmetric in ξ, η, we obtain that

$$[[T_\xi, T_\eta], m_t].$$

The algebra generated by $\left\{m_t : t \in \mathbb{R}^N\right\}$ is Π^N. For any $p \in \Pi^N$ we have $[T_\xi, T_\eta] p = p [T_\xi, T_\eta] 1 = 0$. □

One important consequence of the Theorem 2.7 is that the operators $T_1, \ldots, T_N$ generate a commutative algebra.

3. INTERTWINING OPERATOR IN THE CASE OF ORTHOGONAL SYSTEM

In this section, we give an intertwining operator between T_ξ and the directional derivative ∂_ξ in the case of orthogonal subsystem of root system R. More precisely, we consider a system $R = \{\alpha_1, \dots, \alpha_n\}$ composed by n vectors of $\mathbb{R}^N$, which are pairwise orthogonal and $\kappa = (\kappa_1, \dots, \kappa_n) \in (0, \infty)^n$ and $\xi \in \mathbb{R}^N$. The associated operator T_ξ with R and κ takes the form

$$(T_\xi f)(x) \;=\; \partial_\xi f(x) + \sum_{j=1}^{n} \kappa_j \langle \alpha_j, \xi\rangle \frac{f(x) - f(\tau_{\alpha_j} x)}{\langle x, \alpha_j\rangle}. \tag{3.1}$$

Let $h : \mathbb{R}^n \times \mathbb{R}^N \to \mathbb{R}^N$ be the function defined by

$$h(t, x) = x + \sum_{j=1}^{n} (t_j - 1) \frac{\langle x, \alpha_j\rangle}{|\alpha_j|^2} \alpha_j, \tag{3.2}$$

where $t = (t_1, \dots, t_n) \in \mathbb{R}^n$ and $x \in \mathbb{R}^N$.
We define

$$\chi_\kappa(f)(x) = \frac{1}{\Gamma(\kappa)} \int_{[0,1]^n} f(h(t, x)) w(t) dt, \tag{3.3}$$

where $w(t) = \prod_{j=1}^{n} (1 - t_j)^{\kappa_j - 1}$ and $\Gamma(\kappa) = \prod_{j=1}^{n} \Gamma(\kappa_j)$.

Theorem 3.1. *Let $f \in C^\infty(\mathbb{R})$, then we have*

$$T_\kappa \circ \chi_\kappa f(x) = \chi_\kappa \circ \partial_\xi f(x). \tag{3.4}$$

Proof. For $j = 1, \dots, n$, we denote by θ_j the orthogonal projection in $\mathbb{R}^n$, with respect to the hyperplane $(\mathbb{R}e_j)^\perp$ orthogonal to the vector e_j of the canonical basis $(e_1, \dots, e_n)$ of $\mathbb{R}^n$. The orthogonal projection θ_j acts on $\mathbb{R}^n$ as

$$\theta_j(t) = (t_1, \dots, t_{j-1}, 0, t_{j+1}, \dots, t_n).$$

The system R is orthogonal, then for $j = 1, \dots, n$, we have

$$\begin{aligned} h(t, \tau_{\alpha_j} x) &= \tau_{\alpha_j} x + \sum_{k=1}^{n} (t_k - 1) \frac{\langle \tau_{\alpha_j} x, \alpha_k\rangle}{|\alpha_k|^2} \alpha_k \\ &= x - \frac{\langle x, \alpha_j\rangle}{|\alpha_j|^2} \alpha_j + \sum_{k=1,\, k\neq j}^{n} (t_k - 1) \frac{\langle x, \alpha_k\rangle}{|\alpha_k|^2} \alpha_k \\ &= h(\theta_j t, x). \end{aligned}$$

Let $f \in C^\infty(\mathbb{R})$ and $\xi \in \mathbb{R}^N$. The mapping $x \to h(t, x)$ is linear on $\mathbb{R}^N$, then we can write

$$\begin{aligned} \partial_\xi (f(h(t, x))) &= \partial_{h(t,\xi)} f(h(t, x)) \\ &= \partial_\xi f(h(t, x)) + \sum_{j=1}^{n} (t_j - 1) \frac{\langle \xi, \alpha_j\rangle}{|\alpha_j|^2} \partial_{\alpha_j} f(h(t, x)). \end{aligned}$$

Hence,

$$\begin{aligned}\partial_\xi \chi_\kappa(f)(x) &= \frac{1}{\Gamma(\kappa)}\int_{[0,1]^n} \partial_\xi(f(h(t,x)))w(t)dt\\ &= \frac{1}{\Gamma(\kappa)}\int_{[0,1]^n} \partial_\xi f(h(t,x))w(t)dt\\ &+ \frac{1}{\Gamma(\kappa)}\sum_{j=1}^n \frac{\langle \xi,\alpha_j\rangle}{|\alpha_j|^2}\int_{[0,1]^n}(t_j-1)\partial_{\alpha_j} f(h(t,x))w(t)dt.\end{aligned}$$

Since we can write

$$\partial_{t_j} f(h(t,x)) = \frac{\langle x,\alpha_k\rangle}{|\alpha_k|^2}\partial_{\alpha_j} f(h(t,x))$$

and

$$\int_0^1 (1-t_j)^{\kappa_j}\partial_{t_j} f(h(t,x))dt = -f(h(\theta_j t,x)) + \kappa_j\int_0^1 (1-t_j)^{\kappa_j-1} f(h(t,x))dt$$

we are lead to

$$\begin{aligned}\int_{[0,1]^n}\partial_{\alpha_j} f(h(t,x))(t_j-1)w(t)dt &= \frac{|\alpha_j|^2}{\langle x,\alpha_j\rangle}\int_{[0,1]^n}\partial_{t_j} f(h(t,x))(t_j-1)w(t)dt\\ &= \kappa_j\frac{|\alpha_j|^2}{\langle x,\alpha_j\rangle}\int_{[0,1]^n}(f(h(\theta_j t,x))-f(h(t,x)))w(t)dt\\ &= -\kappa_j\Gamma(\kappa)\frac{|\alpha_j|^2}{\langle x,\alpha_j\rangle}\big(\chi_\kappa(f)(x)-\chi_\kappa(f)(\tau_{\alpha_j}x)\big).\end{aligned}$$

This, combined with the last expression of $\partial_\xi(\chi_\kappa f)(x)$, yields

$$\partial_\xi \chi_\kappa(f)(x) = \chi_\kappa(\partial_\xi f)(x) - \sum_{j=1}^n \kappa_j\langle\xi,\alpha_j\rangle\frac{\chi_\kappa(f)(x)-\chi_\kappa(f)(\tau_j x)}{\langle x,\alpha_j\rangle}.$$

Therefore,

$$T_\xi(\chi_\kappa f)(x) = \chi_\kappa(\partial_\xi f)(x).$$

□

4. INTERESTING CASES

4.1. **The rank one case.** In the one dimensional case there is only one Dunkl-type operator T_κ associated up to scaling and it equals to

$$T_\kappa f(x) = f'(x) + \kappa\frac{f(x)-f(0)}{x}.$$

This operator leaves the space of polynomials invariant and acts on the monomials as

$$T_\kappa 1 = 0,\ T_\kappa x^n = (n+\kappa)x^{n-1},\ n=1,2,\dots.$$

Its square is given by

$$T_\kappa f(x) = f''(x) + \frac{2\kappa}{x}f'(x) + \kappa(\kappa-1)\frac{f(x)-f(0)}{x^2} - \frac{\kappa(\kappa+1)}{x}f'(0).$$

Consider the confluent hypergeometric function [12, 15]

$$M(a,b;z) = \sum_{n=0}^\infty \frac{(a)_n}{(b)_n}\frac{z^n}{n!}, \tag{4.1}$$

where $(a)_n$ is the Pochammer symbol defined by

$$(a)_n = \frac{\Gamma(a+n)}{\Gamma(a)}. \tag{4.2}$$

This is a solution of the confluent hypergeometric differential equation

$$zy''(z) + (b-z)y'(z) = ay(z). \tag{4.3}$$

This function possesses the following Poisson integral representation [15]

$$M(a,b;z) = \frac{\Gamma(b)}{\Gamma(a)\Gamma(b-a)} \int_0^1 t^{a-1}(1-t)^{b-a-1}e^{zt}dt, \; \Re(b) > \Re(a) > 0. \tag{4.4}$$

Theorem 4.1. *For $\lambda \in \mathbb{C}$ and $\kappa > -1$, the problem*

$$T_\kappa f(x) = i\lambda f(x), \;\; f(0) = 1, \tag{4.5}$$

has a unique analytic solution $M_\kappa(i\lambda x)$ given by

$$M_\kappa(i\lambda x) = M(1, \kappa+1; i\lambda x). \tag{4.6}$$

Proof. Searching a solution of (4.5) in the form $f(z) = \sum_{n=0}^\infty a_n x^n$. Replacing in (4.5), we obtain

$$\sum_{n=0}^\infty (n+1+\kappa)a_{n+1}x^n = i\lambda \sum_{n=0}^\infty a_n x^n$$

Thus,

$$a_{n+1} = \frac{i\lambda}{n+1+\kappa}a_n$$

and

$$a_n = \frac{(i\lambda)^n}{(\kappa+1)_n}.$$

□

Remark 4.1. Multiply the equation (4.5) by x and differentiating both sides, we see that a function u of class C^2 on $\mathbb{R}$, is a solution of the equation (4.5), if and only if, it is a solution of the generalized eigenvalue problem

$$xu'' + (\kappa+1)u' = i\lambda(xu' + u).$$

Proposition 4.2. *The function $\mathbf{M}_\kappa(z)$ defined by*

$$\mathbf{M}_\kappa(z) = \frac{M_\kappa(z)}{\Gamma(\kappa+1)} = \sum_{n=0}^\infty \frac{z^n}{\Gamma(\kappa+1+n)}$$

satisfies the following properties :

(i) *$\mathbf{M}_\kappa(z)$ is analytic in κ and z.*
(ii) *$\mathbf{M}_0(z) = e^z$.*
(iii) *For $\Re(\kappa) > 0$, the function $\mathbf{M}_\kappa(z)$, possesses the integral representation*

$$\mathbf{M}_\kappa(z) = \frac{1}{\Gamma(\kappa)} \int_0^1 (1-t)^{\kappa-1}e^{zt}dt.$$

(iv) *For* $\Re(\kappa) > 0$, *we have*

$$\left|\mathbf{M}_\kappa^{(n)}(z)\right| \leq |z|^n e^{\Re(z)},\ n \in \mathbb{N},\ z \in \mathbb{C}.$$

In particular,

$$|\mathbf{M}_\kappa(i\lambda x)| \leq 1,\ \ \lambda,\ x \in \mathbb{R}.$$

(v) *For* $\Re(\kappa) > 0$, *and all* $x \in \mathbb{R}^*$,

$$\lim_{\lambda\to+\infty} \mathbf{M}_\kappa(i\lambda x) = 0.$$

Proof. (i) and (ii) are trivial. (iii) follows from (4.4). For $n \in \mathbb{N}$, we have

$$M_\kappa^{(n)}(z) = \frac{z^n}{\Gamma(\kappa)} \int_0^1 (1-t)^\kappa t^n e^{zt} dt$$

So we find

$$\begin{aligned}|\mathbf{M}_\kappa^{(n)}(z)| &\leq \frac{|z|^n}{\Gamma(\kappa)} \int_0^1 (1-t)^\kappa e^{\Re(z)t} dt \\ &\leq |z|^n e^{\Re(z)}.\end{aligned}$$

This prove (iv). The limit (iv) follows form (iii) and the Riemann-Lebesgue Lemma.

Definition 4.1. We define the Kummer transform on $L^1(\mathbb{R})$ by

$$\forall\, \lambda \in \mathbb{R}, \qquad \mathcal{F}_\kappa(f)(\lambda) = \int_{\mathbb{R}} f(x)\, \mathbf{M}_\kappa(-i\lambda x)(x)\, dx \tag{4.7}$$

In particular, when $\kappa = 0$, the Kummer transform coincides with the classical Fourier transform

$$\mathcal{F}(f)(\lambda) = \int_{\mathbb{R}} f(x)\, e^{-i\lambda x}\, dx.$$

Theorem 4.3. *Let* f *be a function in* $L^1(\mathbb{R})$ *then* $\mathcal{F}_\kappa(f)$ *belongs to* $C_0(\mathbb{R})$, *where* $C_0(\mathbb{R})$ *is the space of continuous functions having 0 as limit in the infinity. Furthermore,*

$$\|\mathcal{F}_\kappa(f)\|_\infty \leq \|f\|_1. \tag{4.8}$$

Proof. It's clear that $\mathcal{F}_\kappa(f)$ is a continuous function on $\mathbb{R}$.
From proposition 4.2, we get for all $x \in \mathbb{R}^*$,

$$\lim_{\lambda\to+\infty} f(x)\mathbf{M}_\kappa(i\lambda x) = 0$$

and

$$|f(x)\mathbf{M}_\kappa(i\lambda x)| \leq |f(x)|.$$

Since f be in $L^1(\mathbb{R})$, we conclude, using the dominated convergence theorem, that $\mathcal{F}_\kappa(f)$ belongs to $C_0(\mathbb{R})$ and

$$\|\mathcal{F}_\kappa(f)\|_\infty \leq \|f\|_1.$$

□

4.1.1. *Intertwining operator.* We denote by $C^\infty(\mathbb{R})$ the space of infinitely differentiable functions f on $\mathbb{R}$, provided with the topology defined by the semi norms

$$\|f\|_{n,a} = \sup_{\substack{0\le k\le n\\ x\in[-a,a]}} |f^{(k)}(x)|,\ a>0,\ \ n\in\mathbb{N}.$$

In the rank one case the intertwining operator (3.3) is given by

$$(\chi_\kappa f)(x) = \frac{1}{\Gamma(\kappa)}\int_0^1 (1-t)^{\kappa-1} f(tx)\,dt. \tag{4.9}$$

This operator is a particular case, when $\gamma=0$ of the so called Erdélyi-Kober fractional integral $K^{\gamma,\delta}$, which is given by

$$(K^{\gamma,\delta}f)(x) = \frac{1}{\Gamma(\delta)}\int_0^1 (1-t)^{\delta-1}t^\gamma f(tx)\,dt,\ \delta>0,\ \gamma\in\mathbb{R}. \tag{4.10}$$

Theorem 4.4. *For a f function of class C^1 on $\mathbb{R}$, $\chi_\kappa f$ is a function of class C^1 on $\mathbb{R}$ and we have*

$$T_\kappa\circ\chi_\kappa = \chi_\kappa\circ\frac{d}{dx}.$$

Proof. By the theorem of derivation under the integral sign, we can prove that $\chi_\kappa(f)$ is of class C^1 on $\mathbb{R}$ and

$$(\frac{d}{dx}\circ\chi_\kappa f)(x) = \frac{1}{\Gamma(\kappa)}\int_0^1 (1-t)^{\kappa-1} t f'(tx)\,dt,\ \ x\in\mathbb{R}.$$

Using the elementary relation

$$(1-t)^{\kappa-1}t = -(1-t)^\kappa + (1-t)^{\kappa-1}$$

and integration by parts formula we can write, for $x\neq 0$

$$\begin{aligned}(\frac{d}{dx}\circ\chi_\kappa f)(x) &= -\frac{1}{\Gamma(\kappa)}\int_0^1 (1-t)^\kappa f'(tx)\,dt + \frac{1}{\Gamma(\kappa)}\int_0^1 (1-t)^{\kappa-1} f'(tx)\,dt\\ &= \frac{1}{\Gamma(\kappa)}\frac{f(0)}{x} - \frac{\kappa}{x}(\chi_\kappa f)(x) + (\chi_\kappa\circ\frac{d}{dx}f)(x).\end{aligned}$$

The result follows from the fact that

$$(\chi_\kappa f)(0) = \frac{1}{\Gamma(\kappa+1)}f(0).$$

□

Theorem 4.5. *The operator χ_κ, $\kappa>0$, is a topological isomorphism from $C^\infty(\mathbb{R})$ onto itself. The inverse operator χ_κ^{-1} is given for all $f\in C^\infty(\mathbb{R})$ by*

$$\chi_\kappa^{-1}f(x) = \prod_{j=1}^n (1+j+x\frac{d}{dx})(K^{\kappa+1,n-\kappa}f)(x)$$

where $n=\lceil\kappa\rceil$ ($\lceil\kappa\rceil$ denotes the ceiling function the smallest integer $\ge\kappa$).

Proof. Let $a>0$ and $f\in C^\infty(\mathbb{R})$. For $x\in[0,a]$, $t\in[0,1]$ and $l\in\mathbb{N}$, we have the following estimate

$$|t^l(1-t)^\beta f^{(l)}(xt)| \le \|f\|_{l,a}\,(1-t)^{\kappa-1}\ \text{ and }\ \int_0^1 (1-t)^{\kappa-1}dt = \frac{1}{\kappa}.$$

By the theorem of derivation under the integral sign, we can prove that

$$\chi_\kappa f \in C^\infty(\mathbb{R}) \text{ and } \|\chi_\kappa(f)\|_{l.a} \leq \frac{1}{\Gamma(\kappa+1)}\|f\|_{l,a}.$$

Then, χ_κ is a linear continuous mapping from $C^\infty(\mathbb{R})$ onto its self. From formula (21) in [1], the operator

$$\prod_{j=1}^{n}(1+j+x\frac{d}{dx}) \circ K^{\kappa+1,n-\kappa}$$

is a left-inverse of χ_κ

$$\prod_{j=1}^{n}(1+j+x\frac{d}{dx}) \circ K^{\kappa+1,n-\kappa} \circ \chi_\kappa f(x) = f(x) \tag{4.11}$$

Let consider the function

$$g(x) = (\chi_\kappa \prod_{j=1}^{n}(1+j+x\frac{d}{dx}) \circ K^{\kappa+1,n-\kappa} f)(x).$$

Using (4.11), we get

$$\begin{aligned}\prod_{j=1}^{n}(1+j+x\frac{d}{dx}) \circ K^{\kappa+1,n-\kappa} g(x) &= (\prod_{j=1}^{n}(1+j+x\frac{d}{dx})K^{\kappa+1,n-\kappa}\chi_\kappa \\ &\quad \circ \prod_{j=1}^{n}(1+j+x\frac{d}{dx} \circ K^{\kappa+1,n-\kappa} f(x) \\ &= \prod_{j=1}^{n}(1+j+x\frac{d}{dx} \circ K^{\kappa+1,n-\kappa} f(x)\end{aligned}$$

that implies the relation

$$(\prod_{j=1}^{n}(1+j+x\frac{d}{dx}) \circ K^{\kappa+1,n-\kappa}(f-g) = 0. \tag{4.12}$$

The functions f and g are in $C^\infty(\mathbb{R})$, then $K^{\kappa+1,n-\kappa}(f-g) \in C^\infty(\mathbb{R})$.
Since, the linear differential equation

$$(\prod_{j=1}^{n}(1+j+x\frac{d}{dx})y(x) = 0,$$

has a unique C^∞-solution, which is equal to $y(x) = 0$, it follows that

$$K^{\kappa+1,n-\kappa}(f-g)(x) = 0.$$

On the other hand, the operator $K^{\kappa+1,n-\kappa}$ has a left-inverse (see, [1]), then $f = g$. This shows that the operator χ_κ has a right-inverse and finish the proof. □

Let $\kappa > 0$, we define the dual intertwining operator ${}^t\chi_\kappa$ on $\mathcal{D}(\mathbb{R})$ ($\mathcal{D}(\mathbb{R})$ is the space of C^∞-functions on $\mathbb{R}$ with compact support) by

$$({}^t\chi_\kappa f)(x) = \frac{1}{\Gamma(\kappa)} \int_{|x|}^{+\infty} (t-|x|)^{\kappa-1}\, t^{-\kappa}\, f(sgn(x)t)\, dt\ , \quad x \in \mathbb{R}^\star \tag{4.13}$$

Proposition 4.6. *The operator* ${}^t\chi_\kappa$ *is a topological automorphism of* $\mathcal{D}(\mathbb{R})$*, and satisfies the transmutation relation:*

$$\int_{\mathbb{R}}(\chi_\kappa f)(x)g(x)\,dx=\int_{\mathbb{R}}f(x)({}^t\chi_\kappa g)(x)\,dx. \tag{4.14}$$

Proof. Let $f\in C^\infty(\mathbb{R})$ and $g\in\mathcal{D}(\mathbb{R})$, we have

$$\begin{aligned}\int_{\mathbb{R}}(\chi_\kappa f)(x)g(x)\,dx&=\frac{1}{\Gamma(\kappa)}\int_0^{+\infty}\int_0^x(x-t)^{\kappa-1}f(t)\,dt\,f(x)x^{-\kappa}\,dx\\&-\frac{1}{\Gamma(\kappa)}\int_0^\infty\int_0^1(x-t)^{\kappa-1}f(-t)\,dt\,g(-x)x^{-\kappa}\,dx.\end{aligned}$$

Using Fubini's theorem and a change of variable, we get

$$\begin{aligned}\int_{\mathbb{R}}(\chi_\kappa f)(x)g(x)\,dx&=\frac{1}{\Gamma(\kappa)}\int_0^{+\infty}\int_t^\infty x^{-\kappa}(x-t)^{\kappa-1}g(x)\,dx\,f(t)\,dt\\&-\frac{1}{\Gamma(\kappa)}\int_0^\infty\int_{-t}^\infty x^{-\kappa}(x+t)^{\kappa-1}g(-x)\,dx\,f(t)\,dt.\end{aligned}$$

Therefore,

$$\begin{aligned}\int_{\mathbb{R}}(\chi_\kappa f)(x)g(x)\,dx&=\frac{1}{\Gamma(\kappa)}\int_0^{+\infty}\int_{|t|}^\infty x^{-\kappa}(x-|t|)^{\kappa-1}g(sign(t)x)\,dx\,f(t)\,dt\\&=\int_{\mathbb{R}}f(t)({}^t\chi_\kappa g)(t)\,dt.\end{aligned}$$

Proposition 4.7. *Let* $\kappa>0$*, the transformation* $\mathcal{F}_\kappa$ *has the following properties:*

(i) *For* $f\in\mathcal{D}(\mathbb{R})$*,* $\mathcal{F}_\kappa(f)$ *satisfies the decomposition*

$$\mathcal{F}_\kappa(f)=\mathcal{F}\circ^t\chi_\kappa(f).$$

(ii) *For* $f\in\mathcal{D}(\mathbb{R})$*, we have*

$$\mathcal{F}_\kappa(T_\kappa f)(\lambda)=i\lambda\mathcal{F}_\kappa(f)(\lambda).$$

4.2. **Multivariable case.** Let $e_1,\dots,e_N$ be the standard basis of $\mathbb{R}^N$. We denote by τ_k (for each k from 1 to N) the orthogonal projection with respect to the hyperplane orthogonal to e_k, that is to say for every $x=(x_1,\dots,x_N)\in\mathbb{R}^N$

$$\tau_k(x)=x-\tfrac{\langle x,e_k\rangle}{|e_k|^2}e_k=(x_1,\dots,x_{k-1},0,x_{k+1},\dots,x_N).$$

Let $\kappa=(\kappa_1,\kappa_2,\dots,\kappa_N)\in\mathbb{C}^N$. The associated Dunkl type operators T_j for $j=1,\dots N$, are given for $x\in\mathbb{R}^N$ by

$$\begin{aligned}T_jf(x)&=\partial_jf(x)+\sum_{l=1}^N\kappa_l\frac{f(x)-f\big(\tau_l(x)\big)}{\langle x,e_l\rangle}\langle e_k,e_l\rangle\\&=\partial_jf(x)+\kappa_j\frac{f(x)-f\big(x_1,\dots,x_{j-1},0,x_{j+1},\dots,x_N\big)}{x_j}.\end{aligned}$$

These operators form a commuting system. The generalized Laplacian associated with T_j is defined in a natural way as

$$\Delta_\kappa=\sum_{j=1}^NT_j^2. \tag{4.15}$$

Proposition 4.8.

$$\begin{aligned}\Delta_\kappa =&\Delta + 2\sum_{j=1}^N \kappa_j x_j^{-1}\partial_j f(x) - \sum_{j=1}^N (\kappa_j^2+\kappa_j)x_j^{-1}\partial_j f(x_1,\dots,x_{j-1},0,x_{j+1},\dots,x_N))\\ &+\sum_{j=1}^N(\kappa_j^2-\kappa_j)x_j^{-2}(f(x)-f(x_1,\dots,x_{j-1},0,x_{j+1},\dots,x_N)).\end{aligned}$$

This operator will play in our context a similar role to that of the Euclidean Laplacian in the classical harmonic analysis. Obviously, the trivial choice of the multiplicity function $\kappa = 0$ reduces our situation to the analysis related to the classical Laplacian Δ.
Let $\kappa = (\kappa_1, \dots, \kappa_N) \in (0,\infty)^N$. For x, $\lambda \in \mathbb{R}^N$, we consider the function $M_\kappa(\lambda, x)$ which is given as the tensor products

$$M_\kappa(\lambda, x) = \prod_{j=1}^N M_{\kappa_j}(i\lambda_j x_j).$$

Theorem 4.9. *For $\lambda = (\lambda_1, \dots, \lambda_N) \in \mathbb{C}^N$, the function $M_\kappa(\lambda, x)$ is the unique analytic solution of the system*

$$T_k u(x) = i\lambda_k u(x),\ u(0) = 1.$$

References

[1] Yury Luchko, Juan J. Trujillo , Caputo–Type Modification of the Erdlyi–Kober Fractional Derivative, Fractional Calculus and Applied Analysis, Volume 19, Number 3 (2007) 249–267.

[2] I. Cherednik, Double affine Hecke algebras, Knizhnik-Zamolodchikov equations, and Macdonald's operators, *Int. Math. Res. Not.* (1992), no. 9, 171–180.

[3] Dunkl, C. F., Reflection groups and orthogonal polynomials on the sphere, Math. Z., 1988, 197 (1): 33–60.

[4] Dunkl, C. F., Differential–difference operators associated to reflection groups, Trans. Amer. Math. Soc., 1989, 311 (1): 167–183.

[5] Dunkl, C. F. and Opdam E. M., Dunkl operators for complex reflection groups, Proc. London Math. Soc. (3) 86 (2003), 70-108.

[6] Dunkl, C. F., Xu Y., Orthogonal polynomials of several variables, Cambridge: Cambridge Univ. Press 2001, 1-390.

[7] Humphreys, J. E., Reflection groups and Coxter groups Cambridge: Cambridge Univ. Press 1990, 1-204.

[8] I. G. Macdonald, *Affine Hecke algebras and orthogonal polynomials*, Cambridge University Press, 2003.

[9] G. J. Heckman, An elementary approach to the hypergeometric shift operators of Opdam, *Invent. Math.* 103 (1991), 341–350.

[10] G. J. Heckman, Dunkl operators, in *Séminaire Bourbaki, Vol. 1996/97* Astérisque 245 (1997), Exp. No. 828, 4, pp. 223–246;

[11] E.M. Opdam, Dunkl operators, Bessel functions and the discriminant of a finite Coxeter group. Compositio Math. 85 (1993), 333–373.

[12] *NIST Handbook of Mathematical Functions*, Cambridge University Press, 2010; http://dlmf.nist.gov.

[13] de Jeu, M.F., The Dunkl transform. Invent. Math. 113 (1993), 147C162.

[14] M. Rösler, Dunkl operators: theory and applications. Orthogonal polynomials and special functions, E.K. Koelink,W. van Assche (eds.), Spring Lecture Notes in Math. 1817 (2003), 93-36 .

[15] G. N. Watson, A Treatise on the Theory of Bessel Functions, Cambridge University Press, Cambridge, England, 1944.

Department of mathematics, King Saudi University, College of Sciences, P. O Box 2455 Riyadh 11451, Saudi Arabia.

E-mail address: `fbouzaffour@ksu.edu.sa`